

Randy Clark, MD  
 Coral Desert Orthopedics  
 1490 E. Foremaster Dr., Ste. 150  
 St. George, UT 84790  
 435-628-9393 phone  
 435-628-9382 fax

**Dr. Randy Clark**  
 Shoulder Open Sub-Pectoral Biceps Tenodesis  
 Physical Therapy Protocol

	<b>Sling Immobilizer</b>	<b>Range of Motion</b>	<b>Exercises</b>	<b>Goals</b>
<b>0-4 weeks</b>  <i>*Start Physical therapy at 2 weeks</i>	Sling at all times	1-full PROM of shoulder and elbow 2-no active elbow flexion or supination	No lifting/pulling/pushing	1-Decreased pain and swelling 2- full passive ROM
<b>4-6 weeks</b>	Remove sling at 4 weeks	1-Pendulums 2- passive ROM only FF to 120, ER to 30 3-No manipulation	1-Scapular stabilization 2-No resistance training or strengthening 3-Deltoid, Rotator Cuff Isometrics 4-no lifting/pushing/pulling	1- passive FF to 120 2- passive ER to 30 3- decreased pain and swelling
<b>6-8 weeks</b>	No sling	1-advance active assist ROM to full active ROM 2- No manipulation	Same as above	Full ROM
<b>8-12 weeks</b>	No sling	Same as above	1- Begin RTC band strengthening program 2- no weights 3- no lifting/pushing/pulling greater than 3 lbs	Full active ROM with normal scapula-humeral rhythm
<b>12-16 weeks</b>	No sling	Full ROM	1-Begin resistance strengthening	full strength
<b>16 weeks +</b>	No sling	Full ROM	1-Advance resistance strengthening 2-Begin Isokinetics 3-Emphasize eccentric strengthening	ability to perform push-ups, pull-ups, sport-specific movements

*Patients will typically follow up post-operatively in clinic with Dr. Bollier at 2,4, 10, and 24 weeks.*

Specific Recommendations /Restrictions /Goals:

---